

A QUALITATIVE TEST OF A UNIFIED MODEL OF SEYFERT GALAXIES WITH *BEPPOSAX*

M. Cappi^{1*}, L. Bassani¹, G. Malaguti¹, G.G.C. Palumbo^{1,2}, M. Dadina³, A. Comastri⁴, G. Di Cocco¹, P. Blanco⁵, D. Dal Fiume¹, A. Fabian⁶, F. Frontera^{1,7}, M. Guainazzi⁸, T. Maccacaro⁹, R. Maiolino¹⁰, G. Matt¹¹, L. Piro¹², M. Trifoglio¹, and N. Zhang¹³

¹ Istituto Te.S.R.E./CNR, Via Gobetti 101, I-40129 Bologna, Italy

² Dipartimento di Astronomia, Università di Bologna, Italy

³ BeppoSAX S.D.C., ASI, Via Corcolle 19, I-00131 Rome, Italy

⁴ Osservatorio di Bologna, Bologna, Italy

⁵ UCSD, San Diego Ca, USA

⁶ Institute of Astronomy, Cambridge University, Madingley Road, Cambridge CB3 0HA, UK

⁷ Dipartimento di Fisica, Università di Ferrara, Via Paradiso 11, I- 44100 Ferrara, Italy

⁸ Astrophysics Division, SCD -ESA, ESTEC, Postbus 299, 2200 AG Noordwijk, The Netherlands

⁹ Osservatorio Astronomico di Brera, Via Brera 28, I20121 Milano, Italy

¹⁰ Osservatorio Astrofisico di Arcetri, Via L.E. Fermi 5, I5015 Firenze, Italy

¹¹ Dipartimento di Fisica, Università di Roma III, Via della Vasca Navale 84, I00146 Roma, Italy

¹² Istituto di Astrofisica Spaziale, Via Del Fosso del Cavaliere, I-001333 Roma, Italy

¹³ University Space Research Association, Huntsville-Al, USA

* e-mail: mcappi@tesre.bo.cnr.it

ABSTRACT

The broad band 0.1-200 keV spectra of a sample of 5 Seyfert 2 galaxies (NGC 7172, Mkn 3, NGC 2110, NGC 4507 and NGC 7674) have been measured within the first year of the *BeppoSAX* Core program. All sources have been detected up to ~ 100 keV and their spectral characteristics derived with good accuracy. Although the results obtained from the detailed analysis of individual sources indicate some “source-by-source” differences, we show in the following that all spectra are consistent, at least qualitatively, with what expected from a “0th-order” version of unified models. Indeed, a simple test on these data indicates that these Seyfert 2 galaxies are on average intrinsically very similar to Seyfert 1 galaxies (i.e., steep at $E \gtrsim 10$ keV) and that the main difference can be ascribed to a different amount of absorbing matter along the line of sight (i.e. different inclinations of our line of sight with respect to a putative molecular torus or different thicknesses of the tori).

1 PREDICTIONS FROM UNIFIED MODELS

The discovery of broad emission lines in the polarized optical spectra of several Seyfert 2 galaxies (Antonucci & Miller 1985, Tran et al. 1992) has provided the basis for a unified model of Seyfert galaxies in which the main discriminating parameter between Seyfert 1 and Seyfert 2 nuclei is the inclination with respect to our line of sight of a supposed obscuring torus surrounding the central source (see Antonucci 1993 for a review). In this scheme, Seyfert 1 galaxies are active nuclei observed nearly perpendicularly to the torus plane (unabsorbed) whereas Seyfert 2 galaxies represent those seen through the torus (absorbed) and, of course, the main prediction is that Seyfert 2 galaxies are *intrinsically* similar to Seyfert

1 galaxies, once the effects of the torus are properly accounted for.

It is now widely recognized that the *intrinsic* high energy spectrum of Seyfert 1 galaxies consists, on average, of a steep power-law continuum ($\Gamma \sim 1.9$ – 2.0 , Nandra & Pounds 1994) with an exponential cut-off typically at energies larger than 150 keV (Zdziarski et al. 1996, Perola et al. 1998). Therefore, one would expect that Seyfert 2 galaxies exhibit similar high-energy properties. X-ray observations (mainly below ~ 20 keV) of Seyfert 2 galaxies, have shown a variety of spectral characteristics not always consistent with a “0th-order” version of unified models (Smith & Done 1996, Cappi et al. 1996, Turner et al. 1998). However, at high energies where the effects of absorption and matter reprocessing are less evident, measurements are sparse for Seyfert 2 galaxies (but see Johnson et al. 1997). It has been therefore difficult to assess the Seyfert 1 nature of the primary spectrum of Seyfert 2 galaxies from these observations.

Based on these general considerations, we have undertaken a program of observations with *BeppoSAX*, aimed at studying the X-ray spectral properties of bright Seyfert 2 galaxies over a broad energy band (up to about 200 keV) and at testing the validity of unified models. As a matter of fact, *BeppoSAX* can provide crucial information on the intrinsic source properties because high energy photons from a few to several keV can pass through the circumnuclear intervening material and can therefore be compared to the “typical” spectrum of Seyfert 1 galaxies. So far 5 objects have been observed within the first AO cycle, namely NGC 7674 (Malaguti et al. 1998a), Mkn 3 (Cappi et al. 1998), NGC 2110 (Malaguti et al. 1998b), NGC 7172 and NGC 4507 (see also Bassani et al. 1998).

2 A SIMPLIFIED QUALITATIVE TEST

A simple, qualitative, test has been performed on our sources by fitting each source of the sample with the same model: a soft power-law component plus a hard, absorbed, power-law component with reflection and associated Fe K line as illustrated in Fig. 1. In the framework of unified models, the soft power-law is attributed to scattered soft X-ray emission from ionized material placed above the molecular torus, while the hard X-ray and reflection components are interpreted as the direct component absorbed by the torus and its reflection from the inner side of the torus, respectively. The intensity of the soft, scattered, component was assumed to be $\sim 2\%$ that of the direct one. The intensity of the reflection component was fixed to $R = 1$ (corresponding to a 2π coverage as viewed from the X-ray source), and the iron line was assumed to be produced through both the reflection and absorption with an equivalent width of ~ 1 keV (with respect to the reflected continuum) and $\sim 500 \times \frac{N_H}{1.23 \times 10^{24}} \text{ eV}$ (with respect to the direct, absorbed, component), as expected from theoretical models (George & Fabian 1991, Leahy & Creighton 1993). The only free parameters were the intensity and photon index of the direct continuum and the absorption column along the line of sight.

The fit results and unfolded spectra obtained from this test are given in Figure 2. The most interesting result is that all sources are well described by a steep, Seyfert 1-like spectrum, with $\Gamma \sim 1.79$ – 1.95 . This is a newly discovered behaviour of Seyfert 2 galaxies for energies up to ~ 100 keV which supports unified models. It is stressed that this result is largely independent on the presence of the steep soft component and on the assumed intensity of the reflection component. This is demonstrated by the fact that the average Seyfert 2 spectrum obtained averaging all the PDS 20–200 keV data (except NGC 7674) can be well fitted ($\chi^2 \sim 1.2$) by a single power-law model with $\Gamma_{20-200 \text{ keV}} = 1.85 \pm 0.05$ and shows no deviation from the power-law up to ~ 150 keV. The observed major differences in the quality of fits (χ^2_{red} ranging from 0.9–1.8) are to be ascribed to two main factors. The first is source by source differences in the Fe K complex which indicates the need of a more dedicated analysis. The second depends upon the different amount of absorbing matter along the line of sight. The less absorbed source is NGC 2110 which also shows less indication for a reflection component and the most absorbed one is NGC 7674, where only the reflection component is observed possibly because the direct component is completely

blocked by a Compton thick molecular torus (with $N_H \gtrsim 10^{25} \text{ cm}^{-2}$, Malaguti et al. 1998a). Intermediate cases are NGC 7172, NGC 4507 and Mkn 3; in the latter, both the direct and reflected components are clearly resolved spectroscopically. Moreover although a detailed measurement in individual sources of the high-energy cutoff is difficult, it appears clear in the data (see Figure 2) that there is no evidence of such a cutoff for energies up to at least $\sim 100\text{--}150 \text{ keV}$.

REFERENCES

- Antonucci, R.R.J., 1993, *ARA&A*, **31**, 473
Antonucci, R.R.J. & Miller, J.S., 1985, *ApJ*, **297**, 621
Bassani, L., et al. 1998, to appear in proceedings of “Dal nano- al tera -eV: tutti i colori degli AGN”, third Italian conference on AGNs, Roma, *Memorie S.A.It.*, astro-ph/9809327
Cappi M., Mihara, T., Matsuoka, et al., 1996, *ApJ*, **456**, 141
Cappi, M., et al. 1998, *A&A*, in press, astro-ph/9902022
George, I.M. & Fabian, A.C., 1991, *MNRAS*, **249**, 352
Johnson, W.N., Zdziarski, A.A., Madejski, G.M., Paciesas, W.S., Steinle, H., & Lin Y-C, 1997, in proceedings of the Fourth Compton Symposium, Ed. D. Dermere, M.S. Steikman and J.D. Kurfess, *AIP*, 283
Leahy D.A. and Creighton J., 1993, *MNRAS*, **263**, 314
Malaguti, G., et al. 1998a, *A&A*, **331**, 519
Malaguti G. et al. 1998b, *A&A*, in press, astro-ph/9901141
Nandra, K. & Pounds, K.A., 1994, *MNRAS*, **268**, 405
Perola, C., et al. 1998, to appear in proceedings of “Dal nano- al tera -eV: tutti i colori degli AGN”, third Italian conference on AGNs, Roma, *Memorie S.A.It*
Smith, D.A., & Done, C., 1996, *MNRAS*, **280**, 355
Tran, H.D., Miller, J.S. & Kay, L.E., 1992, *ApJ*, **397**, 452
Turner, T.J., George, I.M., Nandra, K., & Mushotzky, R.F., 1998, *ApJ*, **493**, 91
Zdziarski, A.A., Johnson, W.N., Poutanen, J., Magdziarz, P., & Gierlinski, M., 1996, in “The Transparent Universe”, proceedings of the 2nd INTEGRAL Workshop, Ed. C. Winkler, T.J.-L. Courvoiser and P. Durouchoux, ESA SP-382, 373




Figure 1: Simple, qualitative, test model used to fit all the sources of the sample in the framework of unified models.


NGC 2110:

$$N_H \sim 5 \times 10^{22} \text{ cm}^{-2}$$

$$\Gamma \sim 1.97$$

$$\chi_{red}^2 \sim 1.16$$

Malaguti et al. 1998, submitted to A&A


 $\Leftrightarrow$
**NGC 7172:**

$$N_H \sim 10^{23} \text{ cm}^{-2}$$

$$\Gamma \sim 1.83$$

$$\chi_{red}^2 \sim 1.4$$

Dadina et al., in prep


 $\Leftrightarrow$
**NGC4507:**

$$N_H \sim 5 \times 10^{23} \text{ cm}^{-2}$$

$$\Gamma \sim 1.83$$

$$\chi_{red}^2 \sim 1.8$$

Bassani et al., in prep


 $\Leftrightarrow$
**Mkn 3:**

$$N_H \sim 1.3 \times 10^{24} \text{ cm}^{-2}$$

$$\Gamma \sim 1.79$$

$$\chi_{red}^2 \sim 0.9$$

Cappi et al., submitted to A&A

 $\Leftrightarrow$
**NGC 7674:**

$$N_H \gtrsim 10^{25} \text{ cm}^{-2}$$

$$\Gamma \sim 1.95$$

$$\chi_{red}^2 \sim 0.9$$

Malaguti et al. 1998a


 $\Leftrightarrow$


Figure 2: Fit results and broad-band unfolded spectra obtained from the qualitative test. See text for a description of the model fitted. N_H increases going from the top (NGC 2110) to the bottom (NGC 7674).